

November 6, 2009
 ICCUB-08-147, KEK-1292, UB-ECM-PF-08/23

Infinite Sequence of Poincaré Group Extensions: Structure and Dynamics

Sotirios Bonanos¹ and Joaquim Gomis^{2,3}

¹Institute of Nuclear Physics, NCSR Demokritos,
 15310 Aghia Paraskevi, Attiki, Greece

²Departament d'Estructura i Constituents de la Matèria and ICCUB, Universitat de
 Barcelona, Diagonal 647, 08028 Barcelona

³High Energy Accelerator Research Organization (KEK),
 Tsukuba, Ibaraki, 305-0801 Japan

e-mail: sbonano@inp.demokritos.gr
 gomis@ecm.ub.es

Abstract

We study the structure and dynamics of the infinite sequence of extensions of the Poincaré algebra whose method of construction was described in a previous paper [1]. We give explicitly the Maurer-Cartan (MC) 1-forms of the extended Lie algebras up to level three. Using these forms and introducing a corresponding set of new dynamical couplings, we construct an invariant Lagrangian, which describes the dynamics of a distribution of charged particles in an external electromagnetic field. At each extension, the distribution is approximated by a set of moments about the world line of its center of mass and the field by its Taylor series expansion about the same line. The equations of motion after the second extensions contain back-reaction terms of the moments on the world line.

1 Introduction

In a recent paper [1] we have studied aspects of the Chevalley-Eilenberg cohomology of the Galilei and Poincaré groups. In particular we have seen that, at degree two, there is an infinite sequence of Lie algebra extensions, beginning with the Galilei or Poincaré algebras. We recall that the extensions found are non-central, since the corresponding generators transform non-trivially under the corresponding normal subalgebra of the unextended algebra.

In this paper we study further the infinite sequence of extensions of the Poincaré algebra. We study the tensor structure of the extensions and their physical interpretations. It is known that the Poincaré group in $d+1$ ($d > 1$) has no central extensions¹ and that it has a non-central antisymmetric tensor extension [8]. Physically, this extension corresponds to the symmetries of a relativistic particle in a constant electromagnetic field and is known as the Maxwell group [9] [10]. A modification of the Poincaré algebra having only two Lorentz generators, that leaves a constant background electromagnetic field invariant, allows an extension with two central charges that can be interpreted as the electric and magnetic charge (BCR algebra [11]). Non-central extensions have also been considered for the diffeomorphism gauge algebra, see for example [12].

Both central and non-central extensions are controlled by Chevalley-Eilenberg cohomology theory, see for example [13]. Here we compute in a systematic, almost algorithmic², way the most general CE cohomology groups at form degree two. As we will see the non-trivial forms belong to different representations of the subgroup of Lorentz transformations of the Poincaré group, in general with mixed symmetries. The Lorentz transformations are a subgroup of the automorphism group and constitute a normal subgroup of each extended group. We can associate to any non-central extension a Young tableau.

As discussed in [1], the first non-central extension of the Poincaré algebra is obtained by calculating all possible non-trivial closed 2-forms of the subgroup of space-time translations. The forms are closed with respect the exterior differential operator d . The complete extended algebra is constructed from the original algebra and the extensions by incorporating their transformation properties under Lorentz transformations, which is equivalent to replacing the exterior differential operator d by the corresponding “covariant” operator $d + M\wedge$, M being the zero-curvature connection associated to the Lorentz generators: $dM + M \wedge M = 0$.

Once we have an extended algebra, we can further extend it by applying the same procedure to an extended set of “translations” which includes all generators except those of the Lorentz subgroup. In this way we obtain new extensions whose generators belong

¹In $1+1$ dimensions there is one central extension [2], which has been used to study several problems of gravity and Moyal quantization, see for example [3, 4, 5]. Also, in spaces R^{2n} and R^{4n} with automorphism groups $U(n)$ and $Sp(n) \times SU(2)$ (Kähler and hyper-Kähler geometries), Galperin et.al. [6, 7] have obtained complex first-level central extensions (a triplet in hyper-Kähler).

²The calculations make use of the first author’s Mathematica package EDC (Exterior Differential Calculus) [14]. The procedure is described in detail in [1].

to higher dimensional representations of the Lorentz group. This procedure does not end resulting in an infinite sequence of groups – extensions of the Poincaré group. In the limit, our procedure formally defines an infinite Lie algebra. However, we cannot prove this result.

We do not have a precise mathematical interpretation for this infinite Lie algebra. We find it intriguing, however, that the generator content of this algebra is organized in levels like the Lorentzian Kac-Moody algebras that are conjectured to be a symmetry of supergravity; see, for example, [15] for the E_{11} approach and [16] for the E_{10} approach.

In order to obtain a possible physical interpretation of this infinite sequence of extensions of the Poincaré group we construct a relativistic particle Lagrangian, invariant under the extended algebra, by using the MC forms. We also introduce tensor coupling “constants” that we consider as new dynamical variables. These tensor couplings are invariant under the extended symmetries.

The form of the equations of motion following from this Lagrangian leads us to the conclusion that the physical system in question is a distribution of charged particles, described collectively as a particle with a set of multipole moments, moving in a fixed background electromagnetic field. The multipoles can be considered as Goldstone bosons. The background field is described by its Taylor series expansion about the world line of the “particle”, higher terms in the series (and higher moments) appearing with every extension. Moreover, new terms in the equation of motion of the “particle” due to back-reaction terms involving the moments appear. These results are obtained by integrating the equations of motion for the coupling fields and plugging the solutions into the equations of motion of the particle coordinates. Once we choose a particular solution, the equations of motion for the particle coordinates imply a spontaneous breaking of the symmetries of the extended algebra.

The organization of the paper is as follows. In section 2 we introduce our notation and conventions and obtain the first level extensions. We then find explicit expressions for the generators of the extended group, construct the Lagrangian, and deduce the transformations of the fields that leave the Lagrangian invariant. Finally, we obtain the equations of motion for all dynamical variables. In sections 3 and 4 we repeat these steps for the second and third extensions. We also give the defining equations for the fourth level extensions. In section 5 we point out that it is advantageous to consider the Young Tableau symmetries of the different extensions; and how symmetry considerations can determine the structure of higher extensions. Finally, in section 6 we compare our results with other approaches for constructing theories with higher symmetry and discuss the implications.

2 The Poincaré group in 3+1 dimensions

The generators of the unextended Poincaré algebra are the translations P_a and the Lorentz transformations M_{ab} , where the tensor indices take the values (0, 1, 2, 3). Denoting by

η_{ab} the Minkowski metric, the algebra is given by³

$$\begin{aligned} [M_{ab}, M_{cd}] &= -i (\eta_{bc} M_{ad} - \eta_{bd} M_{ac} + \eta_{ad} M_{bc} - \eta_{ac} M_{bd}), \\ [P_a, M_{bc}] &= -i (\eta_{ab} P_c - \eta_{ac} P_b). \end{aligned} \quad (2.1)$$

As described in [1], in order to construct the extensions we make use of the left invariant Maurer-Cartan (MC) form, defined by

$$\Omega = -ig^{-1}dg, \quad (2.2)$$

where g represents a general element of the Poincaré group. The MC form satisfies the Maurer-Cartan equation

$$d\Omega + i \Omega \wedge \Omega = 0. \quad (2.3)$$

In components the MC 1-form is written, for a generic Lie algebra, as

$$\Omega = X_A \mathcal{X}^A, \quad (2.4)$$

where X_A are the generators of the Lie algebra satisfying

$$[X_B, X_C] = i f^A_{BC} X_A \quad (2.5)$$

and $\mathcal{X}^A$ are corresponding 1-forms.⁴ Throughout this paper we use the same capital letters in plain and calligraphic font to denote generators and associated 1-forms. The MC equation (2.3) implies that the 1-forms $\mathcal{X}^A$ satisfy

$$d\mathcal{X}^A = \frac{1}{2} f^A_{BC} \mathcal{X}^B \wedge \mathcal{X}^C. \quad (2.6)$$

For the Poincaré case, the MC 1-form (2.4) becomes

$$\Omega = P_a \mathcal{P}^a + \frac{1}{2} M_{ab} \mathcal{M}^{ab}, \quad (2.7)$$

while the MC equation (2.3) in components is

$$\begin{aligned} d\mathcal{P}^a + \mathcal{P}^c \wedge \mathcal{M}_c^a &= 0, \\ d\mathcal{M}^{ab} + \mathcal{M}^{ac} \wedge \mathcal{M}_c^b &= 0. \end{aligned} \quad (2.8)$$

The first step in the cohomological analysis is to freeze the Lorentz degrees of freedom and construct the most general 2-form that can be built from the translations alone, $\mathcal{P}^a$. The MC equations for these generators ($d\mathcal{P}^a = 0$) are obtained by putting $\mathcal{M}^{ab} \rightarrow 0$ in

³ Although the algebra is real and the imaginary units can be made to disappear by replacing all generators G by iG' , we prefer to leave the i 's in the equations because then we can interpret the generators as Hermitian operators.

⁴ In general the generator indices will refer to multiple-index tensors with symmetries. When such indices are summed as in (2.4), (2.5), additional numerical factors must be introduced to compensate for multiple appearances of identical terms in the sum.

(2.8). Then we find that the most general closed invariant 2-form which cannot be written as the differential of an invariant 1-form is

$$\Omega_2 = f_{[ab]} \mathcal{P}^a \wedge \mathcal{P}^b \quad (2.9)$$

where $f_{[ab]}$ is a constant second rank antisymmetric tensor. Therefore the non-trivial 2-form extensions belong to an antisymmetric tensor representation of the Lorentz group. The 1-form “potentials” associated to these closed 2-forms are denoted by $\mathcal{Z}^{[ab]}$, and are defined by the equation

$$d\mathcal{Z}^{[ab]} = \mathcal{P}^a \wedge \mathcal{P}^b. \quad (2.10)$$

From this equation we obtain the algebra of the corresponding generators, denoted by $Z_{[ab]}$. We find

$$[P_a, P_b] = +i Z_{[ab]}, \quad (2.11)$$

which implies that there is no central extension of the Poincaré group.

With the rotations included, the extended set of MC 1-forms satisfies the equations

$$\begin{aligned} d\mathcal{P}^a &= -\mathcal{P}^c \wedge \mathcal{M}_c^a, \\ d\mathcal{M}^{ab} &= -\mathcal{M}^{ac} \wedge \mathcal{M}_c^b, \\ d\mathcal{Z}^{[ab]} &= -\mathcal{Z}^{[ac]} \wedge \mathcal{M}_c^b - \mathcal{M}_c^a \wedge \mathcal{Z}^{[cb]} + \mathcal{P}^a \wedge \mathcal{P}^b. \end{aligned} \quad (2.12)$$

The associated algebra was introduced before in the literature [11, 9, 10]. It is known as the Maxwell algebra.

2.1 Explicit parametrization

Here we will introduce explicit parameters labeling the group elements, which will induce an explicit parametrization of all MC 1-forms in terms of the differentials of these parameters. We will first obtain expressions for the MC 1-forms without the Lorentz generators. Specifically, the general element of this coset of the extended group will be parametrized, locally, by

$$g = e^{iP_a x^a} e^{\frac{i}{2} Z_{ab} \theta^{ab}}. \quad (2.13)$$

where x^a , $\theta^{[ab]}$ are the group parameters associated to the generators P_a , Z_{ab} . The component MC 1-forms (2.4) can be computed directly from the definition (2.2) and the commutator (2.11) using the Baker-Campbell-Hausdorff formula. The result is

$$\mathcal{P}^a = dx^a, \quad \mathcal{Z}^{[ab]} = d\theta^{[ab]} + \frac{1}{2}(x^a dx^b - x^b dx^a). \quad (2.14)$$

If we want to have the explicit expressions of these MC 1-forms when we include the Lorentz degrees of freedom, the right hand side of all vector and tensor expressions given above must be multiplied by an appropriate orthogonal matrix, U , for each index:

$$\mathcal{P}^a = U^{-1a}{}_b dx^b, \quad \mathcal{Z}^{[ab]} = U^{-1a}{}_p U^{-1b}{}_q \left(d\theta^{[pq]} + \frac{1}{2}(x^p dx^q - x^q dx^p) \right). \quad (2.15)$$

U depends on the parameters associated to the Lorentz generators and can be obtained by adding appropriate terms to (2.13). In terms of U the Lorentz MC 1-forms have the explicit representation $\mathcal{M}^a{}_b = U^{-1}{}^a{}_c dU^c{}_b$.

The vectors fields dual to these MC forms (when we freeze the Lorentz degrees of freedom) are:

$$Z_{ab} = -i \frac{\partial}{\partial \theta^{ab}}, \quad (2.16)$$

$$P_a = -i \left(\frac{\partial}{\partial x^a} + \frac{1}{2} x^r \frac{\partial}{\partial \theta^{ar}} \right), \quad (2.17)$$

$$M_{ab} = -i \left(x_a \frac{\partial}{\partial x^b} - x_b \frac{\partial}{\partial x^a} + \theta_a{}^r \frac{\partial}{\partial \theta^{br}} - \theta_b{}^r \frac{\partial}{\partial \theta^{ar}} \right), \quad (2.18)$$

and satisfy the commutators

$$\begin{aligned} [M_{ab}, M_{cd}] &= -i (\eta_{bc} M_{ad} - \eta_{bd} M_{ac} + \eta_{ad} M_{bc} - \eta_{ac} M_{bd}), \\ [P_a, M_{bc}] &= -i (\eta_{ab} P_c - \eta_{ac} P_b), \\ [P_a, P_b] &= +i Z_{ab}, \\ [M_{ab}, Z_{cd}] &= -i (\eta_{bc} Z_{ad} - \eta_{bd} Z_{ac} + \eta_{ad} Z_{bc} - \eta_{ac} Z_{bd}), \\ [P_a, Z_{bc}] &= 0, \\ [Z_{ab}, Z_{cd}] &= 0. \end{aligned} \quad (2.19)$$

2.2 Particle Lagrangian and Noether charges

To obtain a physical interpretation, one possibility is to construct a particle Lagrangian that is invariant under the extended Poincaré group using the non-linear realization method [17] for space-time symmetries, see for example [18]. A diffeomorphism-invariant free particle Lagrangian is $\mathcal{L}_0 = m\sqrt{-\dot{x}_a^2}$ and depends on the translation 1-forms only. A possible Lagrangian including also the first extension 1-forms $\mathcal{Z}^{ab}$ (2.14), is

$$\mathcal{L} = m\sqrt{-\dot{x}_a^2} + \frac{1}{2} f_{ab} \mathcal{Z}^{ab} = m\sqrt{-\dot{x}_a^2} + \frac{1}{2} f_{ab} (\dot{\theta}^{ab} + x^{[a} \dot{x}^{b]}), \quad (2.20)$$

where we have introduced the antisymmetric tensor couplings $f_{ab}(\tau)$ that are considered as new dynamical variables, in addition to the group space coordinates (x^a, θ^{ab}) , which are now also functions of the particle proper time. The physical interpretation of the extra variables θ^{ab}, f_{ab} will be given after the equations of motion have been obtained.

This Lagrangian is invariant under translations⁵

$$\delta_P x^a = \epsilon^a, \quad (2.21)$$

$$\delta_P \theta^{ab} = -\frac{1}{2} (\epsilon^a x^b - \epsilon^b x^a), \quad (2.22)$$

$$\delta_P f_{ab} = 0, \quad (2.23)$$

⁵The generators of these transformations are the right invariant vector fields.

and the non-vanishing shifts

$$\delta_Z \theta^{ab} = \epsilon^{ab}, \quad \delta_Z f_{ab} = 0. \quad (2.24)$$

The Noether charges associated to these symmetries are

$$P_a = p_a - \frac{1}{2} p_{ab} x^b, \quad Z_{ab} = p_{ab}. \quad (2.25)$$

If we compute the Poisson bracket among these generators we find

$$\{P_a, P_b\} = -Z_{ab}, \quad (2.26)$$

i.e., we recover the algebra⁶ (2.11).

In the proper time gauge the Euler-Lagrange equations of motion following from (2.20) are

$$\delta \theta^{ab} \rightarrow \dot{f}_{ab} = 0, \quad (2.27)$$

$$\delta f_{ab} \rightarrow \dot{\theta}^{ab} + \frac{1}{2} (x^a \dot{x}^b - x^b \dot{x}^a) = 0, \quad (2.28)$$

$$\delta x^a \rightarrow -m \ddot{x}_a + f_{ab} \dot{x}^b = 0. \quad (2.29)$$

Equation (2.29), with $f_{ab} = e F_{ab}$, is the Lorentz force equation determining the motion of a particle of mass m and charge e in an electromagnetic field F_{ab} . Note that for this case the equation of motion for f_{ab} , (2.28) does not affect the dynamics of the coordinates. This equation tells us that $\dot{\theta}^{ab}$ is proportional to the ab component of the angular momentum (or magnetic moment) of the particle.⁷ In other words θ^{ab} is a non-local function of the components of the angular momenta of the particle.

Integration of the equation of motion associated to θ gives $f_{ab} = f_{ab}^0 = e F_{ab}^0$. We see that this solution spontaneously breaks Lorentz symmetry. If we substitute this solution in the equation of motion for the variable x , (2.29), we find that it describes the motion of a particle in a constant, fixed EM field with

$$F_{ab} = F_{ab}^0 = \text{constant}. \quad (2.30)$$

It can be obtained from the potential

$$A_a = -\frac{1}{2} F_{ab}^0 x^b. \quad (2.31)$$

⁶The reason for the overall sign difference from the starting algebra is that now the generators are *active* operators.

⁷The terminology refers to the space-space components; the space-time components give a Lorentz-boosted momentum (or dipole moment).

3 Second level extensions

One can obtain further extensions of the Poincaré group which lead to new generators in higher dimensional representations of the Lorentz group. In order to find them we apply the same procedure as in the last section, at every level taking as “translations” all generators of the previous level other than the Lorentz ones, $\mathcal{M}^{ab}$.

For the second extension we take as “translations” the 1-forms

$$\mathcal{P}^a, \mathcal{Z}^{[ab]}. \quad (3.1)$$

The calculation results in 20 closed non-trivial 2-forms which can be written as the components the tensor⁸

$$2 \mathcal{P}^a \wedge \mathcal{Z}^{[bc]} - \mathcal{P}^b \wedge \mathcal{Z}^{[ca]} - \mathcal{P}^c \wedge \mathcal{Z}^{[ab]}. \quad (3.2)$$

Again, introducing the second-level potential 1-form $\mathcal{Y}^{a[bc]}$, with the same symmetries as the above 2-form tensor and unfreezing the Lorentz freedom, we find

$$\begin{aligned} d\mathcal{Y}^{a[bc]} &= -\mathcal{M}^a_s \wedge \mathcal{Y}^{s[bc]} - \mathcal{M}^b_s \wedge \mathcal{Y}^{a[sc]} - \mathcal{M}^c_s \wedge \mathcal{Y}^{a[bs]} \\ &\quad + 2 \mathcal{P}^a \wedge \mathcal{Z}^{[bc]} - \mathcal{P}^b \wedge \mathcal{Z}^{[ca]} - \mathcal{P}^c \wedge \mathcal{Z}^{[ab]}. \end{aligned} \quad (3.3)$$

The corresponding generators $Y_{a[bc]}$ appear in the commutators of the original translations with the first level extensions

$$[P_a, Z_{[bc]}] = 2 i Y_{a[bc]} - i Y_{b[ca]} - i Y_{c[ab]}. \quad (3.4)$$

The complete set of the second extension commutators are:

$$\begin{aligned} [M_{ab}, M_{cd}] &= -i (\eta_{bc} M_{ad} - \eta_{bd} M_{ac} + \eta_{ad} M_{bc} - \eta_{ac} M_{bd}), \\ [P_a, M_{bc}] &= -i (\eta_{ab} P_c - \eta_{ac} P_b), \\ [P_a, P_b] &= i Z_{ab}, \\ [M_{ab}, Z_{cd}] &= -i (\eta_{bc} Z_{ad} - \eta_{bd} Z_{ac} + \eta_{ad} Z_{bc} - \eta_{ac} Z_{bd}), \\ [P_a, Z_{bc}] &= i (2 Y_{abc} - Y_{bca} - Y_{cab}), \\ [Z_{ab}, Z_{cd}] &= 0, \\ [Y_{pab}, M_{cd}] &= -i (\eta_{bc} Y_{pad} - \eta_{bd} Y_{pac} + \eta_{ad} Y_{pbc} - \eta_{ac} Y_{pbd} + \eta_{pc} Y_{dab} - \eta_{pd} Y_{cab}), \\ [Y_{pab}, Z_{cd}] &= 0, \\ [Y_{pab}, P_c] &= 0, \\ [Y_{pab}, Y_{qcd}] &= 0. \end{aligned} \quad (3.5)$$

Note that, at this level, the operators Z_{ab} and Y_{abc} generate an Abelian subgroup.

⁸This tensor is antisymmetric in $[bc]$ and its totally antisymmetric part vanishes. This leads to 4 identities, $\epsilon_{abcd} \mathcal{P}^b \wedge \mathcal{Z}^{[cd]} = 0$, leaving $4 \times 6 - 4 = 20$ independent components.

3.1 Explicit parametrization

Introducing the second extension parameters $\xi^{a[bc]}$ (coordinates in group space) with the symmetries of $\mathcal{Y}^{a[bc]}$, the coset element in the second extension can be written as

$$g = e^{iP_a x^a} e^{\frac{i}{2}Z_{ab}\theta^{ab}} e^{\frac{i}{2}Y_{abc}\xi^{abc}}. \quad (3.6)$$

We can then compute, as before, the corresponding MC forms; the ones associated to the translations and first extension are not modified, while the second extension MC forms are found to be

$$\mathcal{Y}^{abc} = d\xi^{abc} - 2dx^a\theta^{bc} + dx^b\theta^{ca} + dx^c\theta^{ab} + \frac{1}{2}x^a(x^b dx^c - x^c dx^b). \quad (3.7)$$

The differential operators dual to the 1-forms given above provide a representation of the extended algebra (3.5) (in the summations below, differentiations with respect to variables that are zero – θ^{00} , ξ^{011} , etc., – are omitted):

$$Y_{abc} = -i \frac{\partial}{\partial \xi^{abc}}, \quad (3.8)$$

$$Z_{ab} = -i \frac{\partial}{\partial \theta^{ab}}, \quad (3.9)$$

$$P_a = -i \left(\frac{\partial}{\partial x^a} + \frac{1}{2}x^r \frac{\partial}{\partial \theta^{ar}} + \theta^{rs} \frac{\partial}{\partial \xi^{ars}} - \theta^{rs} \frac{\partial}{\partial \xi^{rsa}} - \frac{1}{2}x^r x^s \frac{\partial}{\partial \xi^{rsa}} \right), \quad (3.10)$$

$$\begin{aligned} M_{ab} = & -i \left[x_a \frac{\partial}{\partial x^b} - x_b \frac{\partial}{\partial x^a} + \theta_a^r \frac{\partial}{\partial \theta^{br}} - \theta_b^r \frac{\partial}{\partial \theta^{ar}} \right. \\ & \left. + \frac{1}{2}(\xi_a^{rs} \frac{\partial}{\partial \xi^{brs}} - \xi_b^{rs} \frac{\partial}{\partial \xi^{ars}}) + \xi_a^{rs} \frac{\partial}{\partial \xi^{rsb}} - \xi_b^{rs} \frac{\partial}{\partial \xi^{rsa}} \right]. \end{aligned} \quad (3.11)$$

3.2 Lagrangian associated to the second level extension

When we include the second extension, the particle Lagrangian becomes

$$\mathcal{L} = m\sqrt{-\dot{x}_a^2} + \frac{1}{2}f_{ab}\mathcal{Z}^{ab} + \frac{1}{2}f_{abc}\mathcal{Y}^{abc}, \quad (3.12)$$

where the new tensor couplings $f_{abc}(\tau)$ have the symmetries of $\mathcal{Y}^{abc}$ and, together with the new group space coordinates $\xi^{abc}(\tau)$, are considered as new dynamical variables.

Apart from the ordinary Lorentz transformations, the non-trivial transformations leaving the Lagrangian invariant are

$$\delta_P x^a = \epsilon^a, \quad (3.13)$$

$$\delta_P \theta^{ab} = -\frac{1}{2}(\epsilon^a x^b - \epsilon^b x^a), \quad (3.14)$$

$$\delta_P \xi^{abc} = -x^a(\epsilon^b x^c - \epsilon^c x^b). \quad (3.15)$$

$$\delta_Z \theta^{ab} = \epsilon^{ab}, \quad (3.16)$$

$$\delta_Z \xi^{abc} = 2x^a \epsilon^{bc} - x^b \epsilon^{ca} - x^c \epsilon^{ab}, \quad (3.17)$$

$$\delta_Y \xi^{abc} = \epsilon^{abc}, \quad (3.18)$$

where ϵ^a , ϵ^{ab} , ϵ^{abc} are arbitrary displacements of the group parameters. The conserved quantities are written as

$$Q = \epsilon^a P_a + \frac{1}{2} \epsilon^{ab} Z_{ab} + \frac{1}{2} \epsilon^{abc} Y_{abc}, \quad (3.19)$$

from which we obtain the Noether generators

$$P_a = p_a - \frac{1}{2} p_{ab} x^b + p_{bca} x^b x^c, \quad Z_{ab} = p_{ab} + x^c (2p_{cab} - p_{bca} - p_{abc}), \quad Y_{abc} = p_{abc}. \quad (3.20)$$

The Poisson brackets among these generators reproduce the algebra (3.5) up to a sign.

The Euler-Lagrange equations of motion now take the form

$$\delta \xi^{abc} \rightarrow \dot{f}_{abc} = 0, \quad (3.21)$$

$$\delta \theta^{ab} \rightarrow \dot{f}_{ab} = (-2f_{cab} + f_{abc} + f_{bca}) \dot{x}^c, \quad (3.22)$$

$$\delta f_{abc} \rightarrow \dot{\xi}^{abc} - 2\dot{x}^a \theta^{bc} + \dot{x}^b \theta^{ca} + \dot{x}^c \theta^{ab} + \frac{1}{2} x^a (x^b \dot{x}^c - x^c \dot{x}^b) = 0, \quad (3.23)$$

$$\delta f_{ab} \rightarrow \dot{\theta}^{ab} + \frac{1}{2} (x^a \dot{x}^b - x^b \dot{x}^a) = 0, \quad (3.24)$$

$$\begin{aligned} \delta x^a \rightarrow -m\ddot{x}_a + f_{ab} \dot{x}^b &= -\frac{1}{2} \dot{f}_{ab} x^b + \frac{1}{2} (-2f_{abc} + f_{bca} + f_{cab}) \dot{\theta}^{bc} \\ &\quad - \frac{1}{2} (-2f_{bca} + f_{cab} + f_{abc}) x^b \dot{x}^c. \end{aligned} \quad (3.25)$$

Substituting for $\dot{f}^{ab}$, $\dot{\theta}^{ab}$ from (3.22), (3.24), we find that the RHS of (3.25) vanishes, so that the equation of motion for x^a depends only on f_{ab} :

$$-m\ddot{x}_a + f_{ab} \dot{x}^b = 0. \quad (3.26)$$

If we integrate equations (3.21), (3.22) for f_{abc} and f_{ab} , we get

$$f_{abc} = f_{abc}^0, \quad f_{ab} = (-2f_{cab}^0 + f_{abc}^0 + f_{bca}^0) x^c + f_{ab}^0, \quad f_{...}^0 = \text{const.} \quad (3.27)$$

Note that these solutions break the extended symmetry spontaneously. If we substitute these expressions back in (3.26) we have

$$-m\ddot{x}_a + f_{ab} \dot{x}^b + (-2f_{cab}^0 + f_{abc}^0 + f_{bca}^0) x^c \dot{x}^b = 0. \quad (3.28)$$

The resulting equation of motion describes a particle in a given EM field which is linear in the cartesian coordinates and can be obtained from the potential

$$A_a = F_{cab}^0 x^b x^c - \frac{1}{2} F_{ab}^0 x^b, \quad f_{cab}^0 = e F_{cab}^0. \quad (3.29)$$

What makes this possible is the symmetry properties of the quantities f_{cab}^0 which imply that the field strength f_{ab} (3.27) satisfies $f_{[ab,c]} = 0$.

Note that, as in the previous level, the equation of motion for f_{ab} and f_{abc} do not affect the dynamics of the coordinates. The variable θ^{ab} retains its old interpretation in terms of the magnetic moment of the particle, while, from (3.23) with $\theta^{ab} = 0$, we see that ξ^{abc} is related to the integral of the first moment of the magnetic moment, i.e., the magnetic quadrupole moment (second moment of the current distribution). Thus, it appears that our physical system is a *distribution* of charged particles, described collectively at this level as the motion of a particle with two sets of moments θ^{ab}, ξ^{abc} , moving in a given EM field. The non-locality of the equations determining the moments suggests that the particles also interact among themselves. This will become apparent at the next level where the equation determining x^a will acquire an extra force term proportional to the magnetic moment.

Writing (3.22) as $df_{ab} = (-2f_{cab} + f_{abc} + f_{bca})dx^c$, we can interpret the coefficients f_{abc} as giving the partial derivatives of f_{ab} .

4 Higher level extensions

In this section we will consider explicitly the higher order extensions up to level four. Here, as we will see, a new phenomenon appears: we need more than one tensor to describe the new extensions. Moreover, some of the lower level extensions no longer commute with themselves or with the translations.

The procedure can be continued indefinitely. It will become apparent that, at level n , several new tensor extensions of rank $n + 1$ appear.

4.1 Third extension

At the third level the procedure gives 60 closed 2-forms which can be arranged as the components of two 4th rank tensors with the following symmetries: $\mathcal{S}_1^{(ab)(cd)}$, $\mathcal{S}_2^{[ab][cd]}$ and the additional antisymmetry $\mathcal{S}_1^{(ab)(cd)} = -\mathcal{S}_1^{(cd)(ab)}$ and $\mathcal{S}_2^{[ab][cd]} = -\mathcal{S}_2^{[cd][ab]}$. These 1-form potentials have, respectively, 45 and 15 independent components and satisfy the equations:

$$d\mathcal{S}_1^{abcd} = \mathcal{P}^a \wedge \mathcal{Y}^{cbd} + \mathcal{P}^a \wedge \mathcal{Y}^{dbc} + \mathcal{P}^b \wedge \mathcal{Y}^{cad} + \mathcal{P}^b \wedge \mathcal{Y}^{dac} - \mathcal{P}^c \wedge \mathcal{Y}^{adb} - \mathcal{P}^c \wedge \mathcal{Y}^{bda} - \mathcal{P}^d \wedge \mathcal{Y}^{acb} - \mathcal{P}^d \wedge \mathcal{Y}^{bca}, \quad (4.1)$$

$$d\mathcal{S}_2^{abcd} = 4 \mathcal{Z}^{ab} \wedge \mathcal{Z}^{cd} + \mathcal{P}^a \wedge \mathcal{Y}^{bcd} - \mathcal{P}^b \wedge \mathcal{Y}^{acd} - \mathcal{P}^c \wedge \mathcal{Y}^{dab} + \mathcal{P}^d \wedge \mathcal{Y}^{cab}. \quad (4.2)$$

In the third extension the new non-vanishing commutators are those with a total of four free indices. From (4.1), (4.2) it follows that they satisfy:

$$[Z_{ab}, Z_{cd}] = 4 i S_{abcd}^2, \quad (4.3)$$

$$[P_a, Y_{bcd}] = i (S_{acbd}^1 - S_{adbc}^1) + \frac{i}{3} (2S_{abcd}^2 - S_{acdb}^2 - S_{adbc}^2), \quad (4.4)$$

where the new generators S_{abcd}^1 , S_{abcd}^2 are assumed to have the full symmetries of the corresponding 1-form potentials.

The coset now will be written

$$g = e^{iP_a x^a} e^{\frac{i}{2} Z_{ab} \theta^{ab}} e^{\frac{i}{2} Y_{abc} \xi^{abc}} e^{\frac{i}{8} S_{abcd}^1 \sigma_1^{abcd}} e^{\frac{i}{8} S_{abcd}^2 \sigma_2^{abcd}}, \quad (4.5)$$

where σ_1^{abcd} , σ_2^{abcd} are new scalar parameters having the symmetries of $\mathcal{S}_1^{abcd}$, $\mathcal{S}_2^{abcd}$. After a long calculation we obtain the following explicit expressions for the new MC 1-forms:

$$\begin{aligned} \mathcal{S}_1^{abcd} &= d\sigma_1^{abcd} - (dx^a \xi^{cbd} + dx^a \xi^{dbc} + dx^b \xi^{cad} + dx^b \xi^{dac}) \\ &\quad + dx^c \xi^{adb} + dx^c \xi^{bda} + dx^d \xi^{acb} + dx^d \xi^{bca} + \frac{1}{2} [x^a x^c (xdx)^{bd} + x^b x^d (xdx)^{ac}], \end{aligned} \quad (4.6)$$

$$\begin{aligned} \mathcal{S}_2^{abcd} &= d\sigma_2^{abcd} - (dx^a \xi^{bcd} - dx^b \xi^{acd} - dx^c \xi^{dab} + dx^d \xi^{cab}) \\ &\quad + 2\theta^{ab} (d\theta^{cd} + (xdx)^{cd}) - 2\theta^{cd} (d\theta^{ab} + (xdx)^{ab}), \end{aligned} \quad (4.7)$$

where we have used the notation $(xdx)^{ab} = (x^a dx^b - x^b dx^a)$.

4.2 Third order Lagrangian and equations of motion

With the third order extensions, the particle Lagrangian becomes

$$\mathcal{L} = m\sqrt{-\dot{x}_a^2} + \frac{1}{2} f_{ab} \mathcal{Z}^{ab} + \frac{1}{2} f_{abc} \mathcal{Y}^{abc} + \frac{1}{8} g_{abcd} \mathcal{S}_1^{abcd} + \frac{1}{8} h_{abcd} \mathcal{S}_2^{abcd}, \quad (4.8)$$

where the new tensor couplings $g_{abcd}(\tau)$, $h_{abcd}(\tau)$ have the symmetries of $\mathcal{S}_1^{abcd}$, $\mathcal{S}_2^{abcd}$, respectively, and together with the new group space coordinates σ_1^{abcd} , σ_2^{abcd} are also treated as new dynamical variables.

This Lagrangian is invariant under the transformations found before plus the following ones for the new variables

$$\delta_p \sigma_1^{abcd} = -\frac{3}{2} (\epsilon^a x^b x^c x^d + \epsilon^b x^a x^c x^d - \epsilon^c x^a x^b x^d - \epsilon^d x^a x^b x^c), \quad (4.9)$$

$$\delta_Z \sigma_1^{abcd} = 3(x^a x^d \epsilon^{bc} + x^a x^c \epsilon^{bd} + x^b x^c \epsilon^{ad} + x^b x^d \epsilon^{ac}), \quad (4.10)$$

$$\delta_Y \sigma_1^{abcd} = x^a (\epsilon^{cbd} + \epsilon^{dbc}) + x^b (\epsilon^{cad} + \epsilon^{dac}) - x^c (\epsilon^{adb} + \epsilon^{bda}) - x^d (\epsilon^{acb} + \epsilon^{bca}), \quad (4.11)$$

$$\delta_{S_1} \sigma_1^{abcd} = \epsilon_1^{abcd} \quad (4.12)$$

and

$$\delta_p \sigma_2^{abcd} = (\epsilon^a x^b - \epsilon^b x^a) \theta^{cd} - (\epsilon^c x^d - \epsilon^d x^c) \theta^{ab}, \quad (4.13)$$

$$\delta_Z \sigma_2^{abcd} = \epsilon^{ac} x^b x^d + \epsilon^{bd} x^a x^c - \epsilon^{ad} x^b x^c - \epsilon^{bc} x^a x^d - 2\epsilon^{ab} \theta^{cd} + 2\epsilon^{cd} \theta^{ab}, \quad (4.14)$$

$$\delta_Y \sigma_2^{abcd} = x^a \epsilon^{bcd} - x^b \epsilon^{acd} - x^c \epsilon^{dab} + x^d \epsilon^{cab}, \quad (4.15)$$

$$\delta_{S_2} \sigma_2^{abcd} = \epsilon_2^{abcd}. \quad (4.16)$$

The conserved quantities are written as

$$Q = \epsilon^a P_a + \frac{1}{2} \epsilon^{bc} Z_{bc} + \frac{1}{2} \epsilon^{abc} Y_{abc} + \frac{1}{8} \epsilon_1^{abcd} S_{abcd}^1 + \frac{1}{8} \epsilon_2^{abcd} S_{abcd}^2, \quad (4.17)$$

from which we obtain the Noether generators

$$P_a = p_a - \frac{1}{2} p_{ab} x^b + p_{bca} x^b x^c - \frac{3}{4} p_{abcd}^1 x^b x^c x^d + \frac{1}{2} p_{abcd}^2 x^b \theta^{cd} \quad (4.18)$$

$$Z_{bc} = p_{bc} + x^d (2p_{dbc} - p_{bcd} - p_{cdb}) + \frac{3}{2} (p_{abcd}^1 - p_{acbd}^1) x^a x^d - \frac{1}{2} (p_{abcd}^2 - p_{acbd}^2) x^a x^d - p_{bcd}^2 \theta^{ad}, \quad (4.19)$$

$$Y_{abc} = p_{abc} + \frac{1}{2} (p_{abcd}^1 - p_{acbd}^1) x^d + \frac{1}{3} (2p_{dabc}^2 - p_{dbca}^2 - p_{dcab}^2) x^d, \quad (4.20)$$

$$S_{abcd}^1 = p_{abcd}^1, \quad S_{abcd}^2 = p_{abcd}^2. \quad (4.21)$$

The Euler-Lagrange equations of motion can be reduced to (3.22), (3.23), (3.24) and the following new equations:

$$\delta \sigma_1^{abcd} \rightarrow \dot{g}_{abcd} = 0, \quad (4.22)$$

$$\delta \sigma_2^{abcd} \rightarrow \dot{h}_{abcd} = 0, \quad (4.23)$$

$$\delta g_{abcd} \rightarrow \dot{\sigma}_1^{abcd} = \dot{x}^a (\xi^{cbd} + \xi^{dbc}) + \dot{x}^b (\xi^{cad} + \xi^{dac}) + \dot{x}^c (\xi^{adb} + \xi^{bda}) + \dot{x}^d (\xi^{acb} + \xi^{bca}) - \frac{1}{2} [x^a x^c (x^b \dot{x}^d - x^d \dot{x}^b) + x^b x^d (x^a \dot{x}^c - x^c \dot{x}^a)], \quad (4.24)$$

$$\delta h_{abcd} \rightarrow \dot{\sigma}_2^{abcd} = \dot{x}^a \xi^{bcd} - \dot{x}^b \xi^{acd} - \dot{x}^c \xi^{dab} + \dot{x}^d \xi^{cab} - \theta^{ab} (x^c \dot{x}^d - x^d \dot{x}^c) + \theta^{cd} (x^a \dot{x}^b - x^b \dot{x}^a), \quad (4.25)$$

$$\delta \xi^{abc} \rightarrow \dot{f}_{abc} = -\dot{x}^d (g_{abcd} - g_{acbd}) - \frac{\dot{x}^d}{3} (2h_{dabc} - h_{dbca} - h_{dcab}), \quad (4.26)$$

$$\delta x^a \rightarrow -m \ddot{x}_a + f_{ab} \dot{x}^b = 0, \quad (4.27)$$

where, in reducing (4.25), (4.27), we have used (3.22), (3.23), (3.24). We should remark that, as with (3.25), the RHS of (4.27) is not identically zero, but vanishes because of the other equations of motion. An extra term that vanishes because of (3.24) also appears on the RHS of (3.22).

As in the previous levels, the last terms in (4.25) allow us to relate σ_1^{abcd} to the third order moments (octupole) of the current distribution. However, $\dot{\sigma}_2^{abcd}$ which vanishes when θ and ξ vanish, must be interpreted differently: it arises from non-linear couplings of the current with the quadrupole moment ($\dot{x}^a \xi^{cbd}$ terms) as well as of θ with $\dot{\theta}$ (using (3.24), the last two terms in (4.25) are $2\theta^{ab}\dot{\theta}^{cd} - 2\theta^{cd}\dot{\theta}^{ab}$).

Integrating (4.22), (4.23), (4.26) and substituting in (3.22), we obtain the equation satisfied by f_{ab} :

$$\dot{f}_{ab} = 3(g_{cabd}^0 - g_{cbad}^0) x^c \dot{x}^d - (2h_{cdab}^0 - h_{cabd}^0 - h_{cbda}^0) x^c \dot{x}^d + (-2f_{cab}^0 + f_{abc}^0 + f_{bca}^0) \dot{x}^c. \quad (4.28)$$

Writing $2x^c \dot{x}^d = \frac{d}{dt}(x^c x^d - 2\theta^{cd})$, as follows from (3.24), we can integrate this equation to get:

$$f_{ab} = \frac{3}{4}(g_{cabd}^0 - g_{cbad}^0 + g_{dabc}^0 - g_{dbac}^0)x^c x^d + \frac{1}{4}(h_{cabd}^0 - h_{cbad}^0 + h_{dabc}^0 - h_{dbac}^0)x^c x^d - 2h_{abcd}^0 \theta^{cd} + (-2f_{cab}^0 + f_{abc}^0 + f_{bca}^0)x^c + f_{ab}^0. \quad (4.29)$$

We observe that, when $h_{abcd}^0 \neq 0$, the tensor f_{ab} depends on θ^{ab} in addition to having terms quadratic in the cartesian coordinates. Thus, only part of f_{ab} can be derived from a potential, and the interaction described by f_{ab} can no longer be interpreted as a pure electromagnetic field. The part of f_{ab} that cannot be derived from a potential gives terms to the equation of motion that couple to the magnetic moment. Thus we write (4.27) as

$$m\ddot{x}_a + 2h_{abcd}^0 \dot{x}^b \theta^{cd} - h_{abcd}^0 x^b \dot{\theta}^{cd} = e F_{ab} \dot{x}^b, \quad (4.30)$$

where F_{ab} represents an ordinary external EM field, now quadratic in the coordinates,

$$F_{ab} = \frac{3}{4}(g_{cabd}^0 - g_{cbad}^0 + g_{dabc}^0 - g_{dbac}^0)x^c x^d + (-2f_{cab}^0 + f_{abc}^0 + f_{bca}^0)x^c + F_{ab}^0. \quad (4.31)$$

The symmetries of g_{abcd} implies that $F_{[ab,c]} = 0$ and thus can be derived from a potential. The terms depending on h_{abcd}^0 in the equation of motion imply a damping effect due to the magnetic moment. As the magnetic moment is determined by the position x , this describes a back-reaction altering the time evolution of x . Thus the dynamics of the motion now depends on the dynamics of the new variables θ^{ab} . We can conjecture that at the next level the tensor f_{ab} will contain terms depending on ξ^{abc} , terms cubic in x^a and terms with the product $\theta^{ab} x^c$ and thus the equation of motion for the coordinates will also couple to the quadrupole moment.

As we did with (3.22), writing (4.26) as $df_{abc} = -dx^d(g_{abcd} - g_{acbd}) - dx^d(2h_{dabc} - h_{dbca} - h_{dcab})/3$, we can interpret the coefficients g_{abcd} and h_{abcd} as giving the partial derivatives of f_{abc} , which, in turn, determine the partial derivatives of f_{ab} . Thus g_{abcd} and h_{abcd} are, effectively, the second derivatives of f_{ab} . Of course, the physical meaning of these two types of second derivatives is different. The terms with h_{abcd} , leading to non-zero $f_{[ab,c]}$, can be interpreted as magnetic sources. It is known that Maxwell's theory is consistent with the existence of such sources. However, we prefer to interpret these terms as introducing a coupling to the magnetic moment $\dot{\theta}^{ab}$ in the equation of motion for the position variables.

The form (4.30) of the equation of motion, together with the equations determining the evolution of the different moments ($\theta^{ab}, \xi^{abc}, \sigma^{abcd}, \dots$), reinforce our conclusion, proposed at the end of section 3, that the physical system described here is a distribution of charges moving consistently (including effects due to non-vanishing moments) in an given EM field. The description is approximated by a series expansion of the field and the corresponding collection of moments of the current distribution, successive levels in the extension procedure giving higher approximations.

At the mathematical level, the coordinates of the extended group space describe the degrees of freedom in the multipole expansion of the current distribution and the induced

interactions between them. And the symmetry group describes how changes in the coordinates and multipole moments are interrelated in order that a self-consistent interpretation in terms of moving charges in a given external EM field, including back-reaction terms, be possible.

4.3 Fourth extension

At the 4th level, the procedure gives 204 new extensions which can be grouped as the components of 5 different 5th rank tensors with definite symmetries. The corresponding 1-form generators, denoted by the symbols $\mathcal{T}_i$, $i = 1, \dots, 5$ and having the symmetries indicated, satisfy:

$$d\mathcal{T}_1^{(abcd)e} = \mathcal{P}^{(a} \wedge \mathcal{S}_1^{bcd)e}, \quad (4.32)$$

$$\begin{aligned} d\mathcal{T}_2^{(abc)(de)} &= \mathcal{S}_1^{(abc)d} \wedge \mathcal{P}^e + \mathcal{S}_1^{(abc)e} \wedge \mathcal{P}^d - 4(\mathcal{Z}^{d(a} \wedge \mathcal{Y}^{bc)e} + \mathcal{Z}^{e(a} \wedge \mathcal{Y}^{bc)d}) \\ &\quad + \frac{4}{3}(\mathcal{P}^{(a} \wedge \mathcal{S}_1^{bcd)e} + \mathcal{P}^{(a} \wedge \mathcal{S}_1^{bce)d}), \end{aligned} \quad (4.33)$$

$$d\mathcal{T}_3^{(abc)[de]} = (\mathcal{S}_1^{(abc)d} \wedge \mathcal{P}^e - \mathcal{S}_1^{(abc)e} \wedge \mathcal{P}^d + \frac{8}{5}(\mathcal{Z}^{d(a} \wedge \mathcal{Y}^{bc)e} - \mathcal{Z}^{e(a} \wedge \mathcal{Y}^{bc)d}) \quad (4.34)$$

$$\begin{aligned} &\frac{6}{5}(\mathcal{S}_2^{adbe} \wedge \mathcal{P}^c - \mathcal{S}_2^{aebd} \wedge \mathcal{P}^c)_{(abc)} - \frac{4}{5}(\mathcal{P}^{(a} \wedge \mathcal{S}_1^{bcd)e} - \mathcal{P}^{(a} \wedge \mathcal{S}_1^{bce)d}), \\ d\mathcal{T}_4^{[abc][de]} &= \mathcal{S}_2^{[abc]d} \wedge \mathcal{P}^e - \mathcal{S}_2^{[abc]e} \wedge \mathcal{P}^d + 4(\mathcal{Z}^{d[a} \wedge \mathcal{Y}^{bc]e} - \mathcal{Z}^{e[a} \wedge \mathcal{Y}^{bc]d}) \end{aligned} \quad (4.35)$$

$$\begin{aligned} &+ \frac{4}{3}(\mathcal{P}^{[a} \wedge \mathcal{S}_2^{bcd]e} - \mathcal{P}^{[a} \wedge \mathcal{S}_2^{bce]d} + 2\mathcal{Y}^{e[ab} \wedge \mathcal{Z}^{cd]} - 2\mathcal{Y}^{d[ab} \wedge \mathcal{Z}^{ce]}), \\ d\mathcal{T}_5^{[abcd]e} &= \mathcal{P}^{[a} \wedge \mathcal{S}_2^{bcd]e} + 2\mathcal{Y}^{e[ab} \wedge \mathcal{Z}^{cd]}, \end{aligned} \quad (4.36)$$

where the subscript (abc) indicates the symmetry operation that must be applied to the expression in parenthesis. These 5th rank tensors have, respectively, (84, 60, 36, 20, 4) independent components. We will not investigate further the 4th level extensions.

5 Young tableau symmetries and possible 5th level extensions

To understand the structure of the higher level extensions, it is helpful to discuss their symmetry properties in terms of Young Tableaux (YT). The YT symmetries of all generators (MC 1-forms) up to level 4 are⁹ :

⁹The third level extensions have no particular YT symmetry, but can be expressed in terms of such tensors: $S_1^{(ab)(cd)} = YT_{31}^{(abc)d} + YT_{31}^{(abd)c} - YT_{31}^{(cda)b} - YT_{31}^{(cdb)a}$, $S_2^{[ab][cd]} = YT_{211}^{[abc]d} - YT_{211}^{[abd]c} - YT_{211}^{[cda]b} + YT_{211}^{[cdb]a}$, where the indices on YT tensors indicate the number of boxes in each row of the YT diagram. Similarly, for the fourth extension tensors $\mathcal{T}_i$ defined in subsection 4.3 to have the corresponding YT symmetry certain symmetry operations must be performed on each $\mathcal{T}_i$.

$$\mathcal{P}^a \square, \mathcal{Z}^{ab} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \mathcal{Y}^{abc} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}, \mathcal{S}_1^{abcd} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}, \mathcal{S}_2^{abcd} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array},$$

$$\mathcal{T}_1^{abcde} \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}, \mathcal{T}_2^{abcde} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}, \mathcal{T}_3^{abcde} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}, \mathcal{T}_4^{abcde} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \mathcal{T}_5^{abcde} \begin{array}{|c|} \hline \square \\ \hline \end{array}.$$

Note that completely symmetric YT do not appear, as the requirement that the exterior derivative of these 1-forms be given in terms of the wedge-product of lower generators implies at least one antisymmetry. The 5th level generators (MC 1-forms) will have 6 indices and will transform as the components of 6th rank tensors with the following possible (in 4 dimensions) YT symmetries:

$$\mathcal{W}_1^{abcdef} \begin{array}{|c|c|c|c|} \hline \square & \square & \square & \square \\ \hline \end{array}, \mathcal{W}_2^{abcdef} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}, \mathcal{W}_3^{abcdef} \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}, \mathcal{W}_4^{abcdef} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array},$$

$$\mathcal{W}_5^{abcdef} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \mathcal{W}_6^{abcdef} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \mathcal{W}_7^{abcdef} \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}, \mathcal{W}_8^{abcdef} \begin{array}{|c|} \hline \square \\ \hline \end{array}.$$

The exterior derivative of each $\mathcal{W}_i$ tensor will then be given in terms of wedge products of pairs of lower order generators whose YT can be combined to give the YT of $\mathcal{W}_i$. For example, the YT $\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}$ can be obtained by multiplying the following pairs of YT: $\begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array} \otimes \square, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \otimes \square, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \otimes \begin{array}{|c|} \hline \square \\ \hline \end{array}, \begin{array}{|c|} \hline \square \\ \hline \end{array} \otimes \begin{array}{|c|} \hline \square \\ \hline \end{array}.$

Thus, we expect $d\mathcal{W}_2^{(abcd)(ef)}$ to be given as a linear combination of the following 4 terms: $\mathcal{T}_1^{(abcd)(e)} \wedge \mathcal{P}^f$, $\mathcal{P}^{(a} \wedge \mathcal{T}_2^{bcd)(ef)}$, $\mathcal{Z}^{e(a} \wedge \mathcal{S}_1^{bcd)f} + \mathcal{Z}^{f(a} \wedge \mathcal{S}_1^{bcd)e}$ and the $(abcd)(ef)$ part of $\mathcal{Y}^{abe} \wedge \mathcal{Y}^{cdf}$, which is, however, identically zero. Requiring the exterior derivative of this linear combination to be zero, we determine the unknown coefficients (up to an overall constant factor) and thus the equation defining $\mathcal{W}_2$:

$$d\mathcal{W}_2^{(abcd)(ef)} = 10 \mathcal{T}_1^{(abcd)(e)} \wedge \mathcal{P}^f - 3\mathcal{P}^{(a} \wedge \mathcal{T}_2^{bcd)(ef)} + 3(\mathcal{Z}^{e(a} \wedge \mathcal{S}_1^{bcd)f} + \mathcal{Z}^{f(a} \wedge \mathcal{S}_1^{bcd)e}). \quad (5.1)$$

Not all possible YT symmetries with a given number of indices are present: already at level 3, the symmetry $\begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array}$ does not appear. Conversely, some YT symmetries may appear more than once, i.e., the linear combination of possible 2 forms with a given symmetry may not be unique. Due to computer memory limitations, we have done the 5th-level calculations in 3 dimensions and found that the generator $\mathcal{W}_7$ does not appear, while the generators $\mathcal{W}_3$ and $\mathcal{W}_5$ appear with multiplicities 2 and 3, respectively (the generators $\mathcal{W}_6$ and $\mathcal{W}_8$, being antisymmetric in 4 indices, cannot exist).

We observe that in the Lagrangian, the coupling to the most symmetric generator (say $k_{(abcd)e}$ corresponding to $\mathcal{T}_1^{(abcd)e}$) will only contribute a term depending on the coordinates to f_{ab} . Thus, restricting attention to these most symmetric extensions, f_{ab} will satisfy $f_{[ab,c]} = 0$,¹⁰ so that an interpretation in terms of an ordinary electromagnetic field will be always possible.

We observe that these coefficients associated to the most symmetric extensions are in one to one correspondence with the zero forms used by Vasiliev to describe the “unfolded

¹⁰Consider the Taylor expansion of f_{ab} about the origin and let $q_{[ab](cde)}$ denote the coefficients of the cubic term. When f_{ab} satisfies $f_{[ab,c]} = 0$, these coefficients must satisfy $q_{[abc](de)} = 0$. These conditions imply that the $q_{[ab](cde)}$ coefficients have the same YT symmetry as that of the coefficients $k_{(abcd)e}$, which satisfy $k_{(abcde)} = 0$, and therefore the $k_{(abcd)e}$ coefficients determine the part cubic in the coordinates of an antisymmetric tensor f_{ab} satisfying $f_{[ab,c]} = 0$.

dynamics” of the Maxwell equations [19], see also [20].

6 Summary and discussion

We have studied in detail the structure and the particle dynamics of the infinite sequence of extensions of the Poincaré algebra outlined in [1]. We have seen that the generators of the non-central extensions belong to tensor representations of the Lorentz group of increasingly higher rank. We can associate one or more Young tableaux to every extension. Although we have done the calculations in four dimensions, the extensions found exist in any dimension where their symmetry is allowed. We conjecture that the complete set of extensions constitutes an infinite Lie algebra.

We do not have a precise mathematical interpretation for this infinite algebra, but we note that its generator content is organized in levels like the Lorentzian Kac-Moody algebras, that are conjectured to be a symmetry of supergravity [15], [16]. It is not completely unnatural that there might be some connection between the two structures, despite the fact that the fields in the Lorentzian Kac-Moody algebras at level zero include the graviton. Following this direction we have studied analogies with the representations of the over-extension of the G_2 , G_2^{++11} algebras with respect to A_3 ¹². If we disregard the level zero, at level one there is a vector corresponding to P_a , at level two an antisymmetric two tensor Z_{ab} , at level three a mixed generator that corresponds to our Y_{abc} , but at level four only one object that corresponds to S_{abcd}^2 exists; the generator S_{abcd}^1 does not appear. At level 5 only T_{abcde}^4, T_{abcde}^5 appear. At level 6 there appears only $W_{abcdef}^5, W_{abcdef}^8$. Therefore we can conclude that only some of the extensions of the infinite sequence of Poincaré algebras we found appear also at non-zero levels of G_2^{++} . We do not know if one can find an infinite algebra that encompasses all the possible Poincaré extensions.

In order to understand the physical significance of this infinite sequence of extensions of the Poincaré group, we have constructed an invariant Lagrangian that depends linearly on the extensions by introducing tensor coupling “constants” that we consider as new dynamical variables. The physical system described by this Lagrangian is a distribution of charged particles moving in an external electromagnetic field. The description is approximate: the particles are described collectively by their multipole moments about the world line of their center of mass and the field by its Taylor expansion about the same line. New terms in the approximation series appear with every extension. The multipoles can be considered as Goldstone bosons. The higher extensions give back-reaction terms describing the effect of the moments on the world line.

We think that sequential extensions of space-time groups including odd generators, using the same methods as in this paper, might be useful in constructing theories containing fermions. We hope to address this point in the future.

¹¹ d=5 N=2 pure supergravity using G_2^{++} was studied in [21].

¹²We have used the computer program SimplicLie [22] to study the level structure of the corresponding representations.

Acknowledgments

We acknowledge Adolfo Azcarraga, Andrea Barducci, Roberto Casalbuoni, Jaume Gomis, Machiko Hatsuda, Satoshi Iso, Yoshihisa Kitazawa, Kiyoshi Kamimura, Jerzy Lukierski, Shun'ya Mizoguchi, Teake Nutma, Norisuke Sakai and George Savvidy for discussions. This work has been supported by MCYT FPA 2007-66665, CIRIT GC 2005SGR-00564, Spanish Consolider-Ingenio 2010 Programme CPAN (CSD2007-00042).

We are also grateful to the referees for urging us to provide a physical interpretation for the extended group coordinates.

References

- [1] S. Bonanos and J. Gomis, “A note on the Chevalley-Eilenberg Cohomology for the Galilei and Poincaré Algebras,” *J. Phys. A: Math. Theor.* **42** (2009) 145206 [arXiv:hep-th/0808.2243].
- [2] J.-M. Lévy-Leblond, “Group-Theoretical Foundations of Classical Mechanics: The Lagrangian Gauge Problem,” *Comm. Math. Phys.* **12** (1969) 64.
- [3] V. Aldaya, J. A. de Azcarraga and R. W. Tucker, “Group Cohomology Approach To The Quantum Equivalence Principle,” *J. Geom. Phys.* **3**, 303 (1986).
- [4] M. Gadella, M. A. Martin, L. Nieto and M. del Olmo “The Stratonovich-Weyl correspondence for one dimensional kinematical groups,” *J. Math. Phys.* **32** (1991) 1182.
- [5] D. Cangemi and R. Jackiw, “Gauge Invariant Formulations Of Lineal Gravity,” *Phys. Rev. Lett.* **69** (1992) 233 [arXiv:hep-th/9203056].
- [6] A. S. Galperin, E. A. Ivanov, V. I. Ogievetsky and E. Sokatchev, “Gauge Field Geometry from Complex and Harmonic Analyticities. I. Kähler and Self-Dual Yang-Mills Cases,” *Ann. Phys.* **185** (1988) 1.
- [7] A. S. Galperin, E. A. Ivanov, V. I. Ogievetsky and E. Sokatchev, “Gauge Field Geometry from Complex and Harmonic Analyticities. II. Hyper-Kähler Case,” *Ann. Phys.* **185** (1988) 22.
- [8] A. Galindo “Lie algebra extensions of the Poincaré Algebra,” *J. Math. Phys.* **8** (1967) 768.
- [9] R. Schrader, “The Maxwell Group and the Quantum Theory of Particles in Classical Homogeneous Electromagnetic Fields,” *Fortsch. Phys.* **20** (1972) 701.
- [10] J. Beckers and V. Hussin, “Minimal Electromagnetic Coupling Schemes. Ii. Relativistic And Nonrelativistic Maxwell Groups,” *J. Math. Phys.* **24** (1983) 1295.

- [11] H. Bacry, P. Combe and J. L. Richard, “Group-theoretical analysis of elementary particles in an external electromagnetic field. 1. the relativistic particle in a constant and uniform field,” *Nuovo Cim. A* **67** (1970) 267.
- [12] T. A. Larsson, “Extensions of diffeomorphism and current algebras,” arXiv:math-ph/0002016
- [13] J. A. de Azcarraga and J. M. Izquierdo, “*Lie groups, Lie algebras, cohomology and some applications in physics.*” Cambridge Univ. Press, 1995.
- [14] S. Bonanos, “Exterior Differential Calculus”, Mathematica Enhancement, <http://www.inp.demokritos.gr/~sbonano/EDC/>
- [15] P. C. West, “E(11) and M theory,” *Class. Quant. Grav.* **18** (2001) 4443 [arXiv:hep-th/0104081].
- [16] T. Damour, M. Henneaux and H. Nicolai, “E(10) and a ‘small tension expansion’ of M theory,” *Phys. Rev. Lett.* **89**, 221601 (2002) [arXiv:hep-th/0207267].
T. Damour, M. Henneaux and H. Nicolai, “Cosmological billiards,” *Class. Quant. Grav.* **20**, R145 (2003) [arXiv:hep-th/0212256].
- [17] S. R. Coleman, J. Wess and B. Zumino, “Structure of phenomenological Lagrangians. I,” *Phys. Rev.* **177** (1969) 2239;
C. G. . Callan, S. R. Coleman, J. Wess and B. Zumino, “Structure of phenomenological Lagrangians. II,” *Phys. Rev.* **177** (1969) 2247.
- [18] J. Gomis, K. Kamimura and P. C. West, “The construction of brane and super-brane actions using non-linear realisations,” *Class. Quant. Grav.* **23** (2006) 7369 [arXiv:hep-th/0607057].
- [19] M. A. Vasiliev, *Int. J. Geom. Meth. Mod. Phys.* **3** (2006) 37 [arXiv:hep-th/0504090].
- [20] E. D. Skvortsov, *JHEP* **0807** (2008) 004 [arXiv:hep-th/0801.2268].
- [21] S. Mizoguchi, K. Mohri and Y. Yamada, “Five-dimensional supergravity and hyperbolic Kac-Moody algebra $G(2)(H)$,” *Class. Quant. Grav.* **23** (2006) 3181 [arXiv:hep-th/0512092].
- [22] T. Nutma “SimpLie: a simple program for Lie algebras”, <http://code.google.com/p/simplie/>